Cholacol® II

Cholacol II Promotes Natural Intestinal Cleansing

The entire gastrointestinal (GI) tract is a muscular tube lined with mucous membranes and is approximately 9 meters (29.5 feet) in length. All of the nutrients we ingest are digested and absorbed in this canal. Solids and liquids are taken in through the mouth, and travel down the esophagus and into the stomach where the first stage of digestion takes place. The contents of the stomach are then emptied into the small intestine, the longest portion of the GI tract. The small intestine accounts for nearly 7 meters (23 feet) of the entire 9-meter digestive tract. It is here that further digestion occurs and the majority of food absorption takes place. Material that passes through the small intestine and into the large intestine is then passed out of the body as waste. Bentonite, also known as montmorillonite, is a colloidal, hydrated aluminum silicate with highly adsorptive properties. The bentonite in Cholacol II works like a magnet to help carry waste materials through the intestines for proper elimination. Bile salts contribute further to intestinal cleansing by helping break down fats. Collinsonia root has been used for centuries by the Chinese as a "bowel tonic" to help keep the intestines clean and functioning properly.

How Cholacol II Keeps You Healthy

Promotes intestinal health

The ingredients found in Cholacol II work together to help cleanse the intestines. Collinsonia root helps keep mucous membranes that line the GI tract in good working order. Bile salts contribute to the breakdown of fats in the intestines. Bentonite attracts and helps carry waste materials through the intestines for appropriate elimination.

Encourages regularity

The combined cleansing and emulsifying properties of the ingredients in Cholacol II help keep the intestines free of the debris that can cause some types of irregularity. †



Introduced in 1957



Content:

90 tablets

Suggested Use: Four tablets, 15 minutes before each meal, or as directed.

Supplement Facts:
Serving Size: 4 tablets
Servings per Container: 22

	per Serving	%DV
Calories	2	
Total Carbohydrate	1 g	<1%*
Sugars	1 g	
Iron	1 mg	4%
Sodium	15 mg	<1%

Amount

*Percent Daily Values (DV) are based on a 2,000-calorie diet.

Proprietary Blend: 1,765 mg Bentonite (montmorillonite), collinsonia (root), and purified bovine bile salts.

Other Ingredients: Honey and calcium stearate.

Four tablets supply approximately: 1,660 mg bentonite, 60 mg collinsonia root powder, and 28 mg purified bovine bile salts.

Warning: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

Sold through health care professionals.

Cholacol® II

What Makes Cholacol II Unique

Product Attributes

This exclusive formula combines the adsorbent properties of bentonite, the astringent properties of collinsonia, and the purgative effects of bile salts to stimulate peristalsis

> Makes Cholacol II an effective cleansing product for the bowel[†]

Each tablet supplies 415 mg of bentonite, 15 mg collinsonia root powder, and 7 mg purified bovine bile salts

> All the benefits of whole foods in a convenient form[†]

Manufacturing and Quality-Control Processes Not disassociated into isolated components

The nutrients in Cholacol II are processed to remain intact, complete nutritional compounds

Degreed microbiologists and chemists in our on-site laboratories continually conduct bacterial and analytical tests on raw materials, product batches, and finished products

> Ensures consistent quality and safety

Vitamin and mineral analyses validate product content and specifications

> Assures high-quality essential nutrients are delivered

Whole Food Philosophy

Our founder, Dr. Royal Lee, challenged common scientific beliefs by choosing a holistic approach of providing nutrients through whole foods. His goal was to provide nutrients as they are found in nature—in a whole food state where he believed their natural potency and efficacy would be realized. Dr. Lee believed that when nutrients remain intact and are not split from their natural associated synergists-known and unknown—bioactivity is markedly enhanced over isolated nutrients. Following this philosophy, even a small amount of a whole food concentrate will offer enhanced nutritional support, compared to an isolated or fractionated vitamin. Therefore, one should examine the source of nutrients rather than looking at the quantities of individual nutrients on product labels.

Studies on nutrients generally use large doses and these studies, some of which are cited below, are the basis for much of the information we provide you in this publication about whole food ingredients. See the supplement facts for Cholacol® II.

Anderson L.E. 1998. Mosby's Medical, Nursing, & Allied Health Dictionary. 5th ed. St. Louis, MO: Mosby: 183, 190, 366, 389, 489,

Balch J.F., Balch P.A. 1997. Prescription for Nutritional Healing. 2nd ed. Garden City Park, NY: Avery Publishing Group: 211-214.

Croce E., et al. 1993. [Rationale for the use of bile salts after cholecystectomy: results of a controlled clinical study using tauroursodeoxycholic acid. Annals of Italian Chiropractic 64(5):

Donovan J.M., Jackson A.A. 1997. Transbilayer movement of fully ionized taurine-conjugated bile salts depends upon bile salt concentration, hydrophobicity, and membrane cholesterol content. *Biochemistry* 36(38): 11444-11451.

Duke J.A., Foster S. 1990. Medical Plants (Eastern/Central), Peterson Field Guides. Boston, MA: Houghton Mifflin: 112-113. Dvorak M. 1989. Ability of bentonite and natural zeolite to adsorb aflatoxin

from liquid media. Veterinary Medicine (Praha) 34(5): 307-316. Frawley D., Lad V. 1992. *The Yoga of Herbs*. 3rd ed. Twin Lakes, WI: Lotus Press: 215. Hoffmann D.L. 1995. *The New Holistic Herbal*. Britain: Barnes & Noble

(special edition by arrangement with Elemental Book Limited); 235. Ivan M., et al. 1992. Effects of bentonite and monensin on selected elements in the stomach and liver of fauna-free and faunated sheep. *Journal of* Dairy Science 75(1): 201-208.

Lewis W.H., Elvin-Lewis P.F. 1977. Medical Botany: Plants affecting Man's Health. New York, NY: John Wiley & Sons: 312-313. Marks J.G., et al. 1995. Prevention of poison ivy and poison oak allergic

contact dermatitis by quaternium-18 bentonite. Journal of the American

Academy of Dermatology 33(2 Pt 1): 212-216.

Pitchford P. 1993. Healing With Whole Foods. Revised ed. Berkeley, CA:
North Atlantic Books: 343-346.

Schwarz T., Werner E. 1990. The effect of long-term bentonite administration on the metabolism of selected trace elements (Fe, Cu, Zn, Mn) in the dwarf goat. Archives of Experimental Veterinary Medicine

Stryock T.R., et al. 1994. Effect of bentonite incorporated in a feed ration with tilmicosin in the prevention of induced Mycoplasma gallisepticu airsacculitis in broiler chickens. Avian Disorders 38(3): 501-505. Stevens J.F., et al. 1999. A novel 2-hydroxyflavone from Collinsonia

canadensis. J Nat Prod 62(2): 392-394.
Sun W. 1994. Studies on isolation and purification of pericillin acylase by adsorption on bentonite. Chinese Journal of Biotechnology 10(2):

Tierra M. 1988. Planetary Herbology. Twin Lakes, WI: Lotus Press

Winston D. 1999. Herbal Therapeutics: Specific Indications For Herbs & Herbal Formulas. 6th ed. Herbal Therapeutics Research Library: 37.

